

Mycophenolate Mofetil Oral Suspension

1g/5ml

Instructions for administration via nasogastric (NG) or percutaneous endoscopic gastrostomy (PEG) tubes:

Ensure that the enteral feeding tube is free from obstruction before administration.

1. Administer the required dose of Mycophenolate Oral Suspension gently and slowly into enteral tube, with a suitable measuring device.

2. Flush the enteral tube with water. A flush volume of 7.5mL is required. However, for longer tubes (over 40cm) flush volume of 50mL should be used.

This product has not been tested with latex NG or PEG tubes and therefore should not be used with tubes made from latex.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Please consult the SmPC for full prescribing information and administration details.



ROS000039-013 February 2024

Rosemont Pharmaceuticals Ltd. Rosemont House, Yorkdale Industrial Park, Braithwaite Street, Leeds LS11 9XE **T** +44 (0)113 244 1400
E infodesk@rosemontpharma.com Sales/ Customer Service: **T** +44 (0) 113 244 1999 **W** www.rosemontpharma.com

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard Adverse events should also be reported to Rosemont Pharmaceuticals Ltd on 0113 244 1400.